



Thrombolytic Checklist

Mark all of the following conditions that currently apply to the patient. Items in italics are for hospital personnel to ascertain. The original of this form should accompany the patient throughout the course of their treatment, and the duplicate copy should be attached to the white copy (original) of the PCR.

ABSOLUTE CONTRAINDICATIONS

- Prolonged CPR
- Head injury
- Blood pressure > 185/110 mm Hg (after treatment with NTG paste)
- Prior cerebral bleeding
- Trauma or surgery within 2 weeks or puncture of noncompressable vessel or organ biopsy
- Central nervous system neoplasm, AV malformation, or aneurysm
- CNS procedure or CVA within 2 months
- Thrombolytic allergy (Streptokinase or anistreplase)
- Active internal bleeding
- Hemorrhagic retinopathy*
- Aortic dissection*

RELATIVE CONTRAINDICATIONS

- Age over 75
- Trauma / surgery more than 2 weeks ago
- Pregnancy or early postpartum
- Chronic severe hypertension
- Recent GI bleeding or active peptic ulcer disease (<10 days)
- History of cerebrovascular accident
- Menstruation
- Anticoagulant medication
- Liver dysfunction
- Terminal cancer or other end-stage disease
- Conditions associated with bleeding risks, such as diabetic retinopathy
- Acute pericarditis or subacute bacterial endocarditis*
- Recent streptococcal infection, if streptokinase or anistreplase will be used*
- Previous Streptokinase / Alteplase*
- Bleeding diathesis*
- Septic thrombophlebitis*